

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Biotinylated Anti-Human 4-1BB Receptor Antibody, Goat, Polyclonal ABT-ABG10004-U025

| | |
|--------------------------|---|
| Artikelname | Biotinylated Anti-Human 4-1BB Receptor Antibody, Goat, Polyclonal |
| Artikelnummer | ABT-ABG10004-U025 |
| Hersteller Artikelnummer | ABG10004-U025 |
| Alternativnummer | ABT-ABG10004-U025-25UG |
| Hersteller | Abcepta |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | ELISA, WB |
| Spezies Reaktivität | Human |
| Klonalität | Polyclonal |
| Reinheit | Produced from sera of goats pre-immunized with highly pure (>98%) recombinant h4-1BB Receptor. Anti-Human 4-1BB Receptor specific antibody was purified by affinity chromatography and then biotinylated. |
| Formulierung | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2. |
| Antibody Type | Polyclonal Antibody |

Anwendungsbeschreibung

WesternBlot: To detect h4-1BB Receptor by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant h4-1BB Receptor is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect h4-1BB Receptor by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.25 - 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems Polyclonal Anti-Human 4-1BB Receptor (60-069P) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant h4-1BB Receptor.. Direct: To detect h4-1BB Receptor by direct ELISA (using 100 µl/well antibody solution) a concentration of 0.25 - 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant h4-1BB Receptor.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.