

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Human EMAP-II Antibody, Goat, Polyclonal ABT-ABG10074-U100

Artikelname	Anti-Human EMAP-II Antibody, Goat, Polyclonal
Artikelnummer	ABT-ABG10074-U100
Hersteller Artikelnummer	ABG10074-U100
Alternativnummer	ABT-ABG10074-U100-100UG
Hersteller	Abcepta
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA, WB
Spezies Reaktivität	Human
Klonalität	Polyclonal
Reinheit	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hEMAP-II. Anti-Human EMAP-II specific antibody was purified by affinity chromatography employing immobilized hEMAP-II matrix.
Formulierung	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Anwendungsbeschreibung

WesternBlot: To detect hEMAP-II by Western Blot analysis this antibody can be used at a concentration of 0.1- 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hEMAP-II is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hEMAP-II by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human EMAP-II (60-401BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hEMAP-II.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.