

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Human Eotaxin Antibody, Goat, Polyclonal ABT-ABG10080-U050

|                          |   |
|--------------------------|---|
| Artikelname              | Anti-Human Eotaxin Antibody, Goat, Polyclonal   |
| Artikelnummer            | ABT-ABG10080-U050   |
| Hersteller Artikelnummer | ABG10080-U050   |
| Alternativnummer         | ABT-ABG10080-U050-50UG  |
| Hersteller               | Abcepta   |
| Wirt                     | Goat  |
| Kategorie                | Antikörper  |
| Applikation              | ELISA, WB   |
| Spezies Reaktivität      | Human   |
| Klonalität               | Polyclonal  |
| Reinheit                 | Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hEotaxin. Anti-Human Eotaxin specific antibody was purified by affinity chromatography employing immobilized hEotaxin matrix. |
| Formulierung             | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.  |
| Antibody Type            | Polyclonal Antibody   |

Anwendungsbeschreibung

WesternBlot: To detect hEotaxin by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hEotaxin is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hEotaxin by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human Eotaxin (60-122GBT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hEotaxin.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.