

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Human Eotaxin-2 Antibody, Goat, Polyclonal ABT-ABG10087-U050

Artikelname	Anti-Human Eotaxin-2 Antibody, Goat, Polyclonal
Artikelnummer	ABT-ABG10087-U050
Hersteller Artikelnummer	ABG10087-U050
Alternativnummer	ABT-ABG10087-U050-50UG
Hersteller	Abcepta
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA, WB
Spezies Reaktivität	Human
Klonalität	Polyclonal
Reinheit	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hEotaxin-2. Anti-Human Eotaxin-2 specific antibody was purified by affinity chromatography employing immobilized hEotaxin-2 matrix.
Formulierung	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Anwendungsbeschreibung

WesternBlot: To detect hEotaxin-2 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hEotaxin-2 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hEotaxin-2 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human Eotaxin-2 (60-123BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hEotaxin-2.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.