

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Human FGF-basic Antibody, Mouse, Monoclonal ABT-ABG10116-U500

Artikelname	Anti-Human FGF-basic Antibody, Mouse, Monoclonal
Artikelnummer	ABT-ABG10116-U500
Hersteller Artikelnummer	ABG10116-U500
Alternativnummer	ABT-ABG10116-U500-500UG
Hersteller	Abcepta
Wirt	Mouse
Kategorie	Antikörper
Applikation	ELISA, IHC, WB
Spezies Reaktivität	Human
Klonalität	Monoclonal
Reinheit	Produced in BALB/c x ICR F1 mice using highly pure recombinant human FGF-basic as the immunizing antigen. This IgG1K antibody was purified from ascites fluid by Protein A affinity chromatography.
Formulierung	A sterile filtered antibody solution was lyophilized from PBS.
Antibody Type	Monoclonal Antibody

Anwendungsbeschreibung

WesternBlot: To detect Human FGF-basic by Western Blot analysis this antibody can be used at a concentration of 0.25-0.50 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Human FGF-basic is 1.0-2.0 ng/lane, under non-reducing conditions.. Sandwich: In a sandwich ELISA (assuming 100µl/well), a concentration of 2.0-4.0 µg/ml of this antibody will detect at least 1000 pg/ml of recombinant human FGF-basic when used with BioGems biotinylated antigen affinity purified anti-human FGF-basic (60-133BT) as the detection antibody at a concentration of approximately 0.125-0.25 µg/ml.. Immunohistochemistry: This antibody stained formalin-fixed, paraffin-embedded sections of human breast invasive ductal carcinoma. The recommended concentrations are 0.25-0.50 µg/ml overnight at 4°C. An HRP-labeled polymer detection system was used with DAB chromogen. Heat induced antigen retrieval was performed with a pH 6.0 Sodium Citrate buffer. Optimal concentrations and conditions may vary. Tissue samples were provided by the Cooperative Human Tissue Network which is funded by the National Cancer Institute.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.