

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Human HCC-1 Antibody, Rabbit, Polyclonal ABT-ABG10153-U050

Artikelname	Anti-Human HCC-1 Antibody, Rabbit, Polyclonal
Artikelnummer	ABT-ABG10153-U050
Hersteller Artikelnummer	ABG10153-U050
Alternativnummer	ABT-ABG10153-U050-50UG
Hersteller	Abcepta
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA, IHC, WB
Spezies Reaktivität	Human
Klonalität	Polyclonal
Reinheit	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hHCC-1. Anti-Human HCC-1 specific antibody was purified by affinity chromatography employing immobilized hHCC-1 matrix.
Formulierung	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Anwendungsbeschreibung

WesternBlot: To detect hHCC-1 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hHCC-1 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hHCC-1 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human HCC-1 (60-174BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hHCC-1.. Immunohistochemistry: This antibody stained formalin-fixed, paraffin-embedded sections of human colon/rectum adenocarcinoma. The recommended concentration is 1.5 mg/ml with an overnight incubation at 4C. An HRP-labeled polymer detection system was used with a DAB chromogen. Proteinase K antigen retrieval is recommended. Optimal concentrations and conditions may vary.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.