

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Human IGF-II Antibody, Rabbit, Polyclonal ABT-ABG10186-U100

Artikelname	Anti-Human IGF-II Antibody, Rabbit, Polyclonal
Artikelnummer	ABT-ABG10186-U100
Hersteller Artikelnummer	ABG10186-U100
Alternativnummer	ABT-ABG10186-U100-100UG
Hersteller	Abcepta
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA, IHC, WB
Spezies Reaktivität	Human
Klonalität	Polyclonal
Reinheit	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hIGF-II (human IGF-II ). Anti-Human IGF-II specific antibody was purified by affinity chromatography employing immobilized hhIGF-II matrix.
Formulierung	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Anwendungsbeschreibung

WesternBlot: To detect hIGF-II by Western Blot analysis this antibody can be used at a concentration of 0.1- 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIGF-II is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hIGF-II by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human IGF-II (60-187BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIGF-II.. Immunohistochemistry: This antibody stained formalin-fixed, paraffin-embedded sections of normal human placenta. The recommended concentration ranges from 0.5 µg/ml-5.0 µg/ml. An HRP-labeled polymer detection system was used with DAB chromogen. Recommended antigen retrievals are heat induced antigen retrieval with a pH 6.0 Sodium Citrate buffer or enzyme induced antigen retrieval with Proteinase K. Optimal concentrations and conditions may vary.. Neutralization: To yield one-half maximal inhibition [ND<span style=font-size: 16px,>50] of the biological activity of hIGF-II (10 ng/ml), a concentration of 3.1 - 4.6 µg/ml of this antibody is required.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.