

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Human IL-12 Antibody, Goat, Polyclonal ABT-ABG10200-U050

Artikelname	Anti-Human IL-12 Antibody, Goat, Polyclonal
Artikelnummer	ABT-ABG10200-U050
Hersteller Artikelnummer	ABG10200-U050
Alternativnummer	ABT-ABG10200-U050-50UG
Hersteller	Abcepta
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA, WB
Spezies Reaktivität	Human
Klonalität	Polyclonal
Reinheit	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hIL12. Anti-Human IL-12 specific antibody was purified by affinity chromatography employing immobilized hIL12 matrix.
Formulierung	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Anwendungsbeschreibung	<p>WesternBlot: To detect hIL-12 by Western Blot analysis this antibody can be used at a concentration of 0.1- 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-12 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hIL-12 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human IL-12 (60-012BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIL-12.. Neutralization: To yield one-half maximal inhibition [ND&lt;span style=font-size: 16px,&gt;50] of the biological activity of hIL-12 (0.3 ng/ml), a concentration of 0.012-0.02 µg/ml of this antibody is required.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.</p>
------------------------	---