

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Human IL-12 Antibody, Mouse, Monoclonal ABT-ABG10204-U500

Artikelname	Anti-Human IL-12 Antibody, Mouse, Monoclonal
Artikelnummer	ABT-ABG10204-U500
Hersteller Artikelnummer	ABG10204-U500
Alternativnummer	ABT-ABG10204-U500-500UG
Hersteller	Abcepta
Wirt	Mouse
Kategorie	Antikörper
Applikation	ELISA, IHC, WB
Spezies Reaktivität	Human
Klonalität	Monoclonal
Reinheit	Produced in BALB/c x ICR F1 mice using highly pure (>98%) recombinant human IL-12 as the immunizing antigen. This IgG1K antibody was purified from ascites fluid by Protein A affinity chromatography.
Formulierung	A sterile filtered antibody solution was lyophilized from PBS.
Antibody Type	Monoclonal Antibody

Anwendungsbeschreibung

WesternBlot: To detect hIL-12 by Western Blot analysis this antibody can be used at a concentration of 0.50-1.0 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-12 is 0.5-1.0 ng/lane, under non-reducing conditions.. Sandwich: In a sandwich ELISA (assuming 100µl/well), a concentration of 2.0-4.0 µg/ml of this antibody will detect at least 1000 pg/ml of recombinant human IL-12 when used with BioGems biotinylated antigen affinity purified anti-human IL-12 (60-012BT) as the detection antibody at a concentration of approximately 4.0-8.0 µg/ml.. Immunohistochemistry: This antibody stained PBMC. The primary antibody was incubated at 10.0 µg/ml for one hour at room temperature followed by a fluorescent labeled secondary antibody. Optimal concentrations and conditions may vary.
 Protocol and staining provided by Dr. Lauren Binge, Laboratory of Prof. Charles Mackay, Monash Univeristy, Australia. <p class=MsoNormal>. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.