

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Human IL-3 Antibody, Rabbit, Polyclonal ABT-ABG10268-U050

|                          |  |
|--------------------------|--|
| Artikelname              | Anti-Human IL-3 Antibody, Rabbit, Polyclonal   |
| Artikelnummer            | ABT-ABG10268-U050  |
| Hersteller Artikelnummer | ABG10268-U050  |
| Alternativnummer         | ABT-ABG10268-U050-50UG   |
| Hersteller               | Abcepta  |
| Wirt                     | Rabbit   |
| Kategorie                | Antikörper   |
| Applikation              | ELISA, WB  |
| Spezies Reaktivität      | Human  |
| Klonalität               | Polyclonal   |
| Reinheit                 | Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hIL-3. Anti-Human IL-3 specific antibody was purified by affinity chromatography employing immobilized hIL-3 matrix. |
| Formulierung             | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.   |
| Antibody Type            | Polyclonal Antibody  |

Anwendungsbeschreibung

WesternBlot: To detect hIL-3 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-3 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hIL-3 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human IL-3 (60-003BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIL-3.. Neutralization: To yield one-half maximal inhibition [ND<span style=font-size: 16px,>50] of the biological activity of hIL-3 (1.0 ng/ml), a concentration of 0.38 - 0.57 µg/ml of this antibody is required.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.