

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Human IL-5 Antibody, Rabbit, Polyclonal ABT-ABG10286-U050

Artikelname	Anti-Human IL-5 Antibody, Rabbit, Polyclonal
Artikelnummer	ABT-ABG10286-U050
Hersteller Artikelnummer	ABG10286-U050
Alternativnummer	ABT-ABG10286-U050-50UG
Hersteller	Abcepta
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA, IHC, WB
Spezies Reaktivität	Human
Klonalität	Polyclonal
Reinheit	Produced from sera of rabbits pre-immunized with highly pure recombinant Human IL-5. Anti-Human IL-5 specific antibody was purified by affinity chromatography employing immobilized Human IL-5 matrix.
Formulierung	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Anwendungsbeschreibung

WesternBlot: To detect Human IL-5 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Human IL-5 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect Human IL-5 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human IL-5 (60-005BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human IL-5.. Immunohistochemistry: This antibody stained formalin-fixed, paraffin-embedded sections of human colon/rectum adenocarcinoma. The recommended concentration is 0.25mg/ml with an overnight incubation at 4C. An HRP-labeled polymer detection system was used with a DAB chromogen. Optimal results for these conditions were achieved without antigen retrieval. Optimal concentrations and conditions may vary.. Neutralization: To yield one-half maximal inhibition [ND<span style=font-size: 16px,>50] of the biological activity of Human IL-5 (5.0 ng/ml), a concentration of 0.05 - 0.15 µg/ml of this antibody is required.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.