

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Murine IL-6 Antibody, Rabbit, Polyclonal ABT-ABG10292-U050

Artikelname	Anti-Murine IL-6 Antibody, Rabbit, Polyclonal
Artikelnummer	ABT-ABG10292-U050
Hersteller Artikelnummer	ABG10292-U050
Alternativnummer	ABT-ABG10292-U050-50UG
Hersteller	Abcepta
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA, IHC, WB
Spezies Reaktivität	Mouse
Klonalität	Polyclonal
Reinheit	Produced from sera of rabbits pre-immunized with highly pure recombinant Murine IL-6. Anti-Murine IL-6 specific antibody was purified by affinity chromatography employing immobilized Murine IL-6 matrix.
Formulierung	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Anwendungsbeschreibung

WesternBlot: To detect Murine IL-6 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Murine IL-6 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect Murine IL-6 by sandwich ELISA (using 100µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Murine IL-6 (61-006BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Murine IL-6.. Immunohistochemistry: This antibody stained colchicine injected mouse brain (cingulate cortex) tissue. The primary antibody was incubated at 0.5 µg/ml overnight at 4°C. This was followed by a peroxidase conjugated secondary antibody and then a fluorescein Tyramide Signal Amplification (TSA(TM)) reagent. Optimal concentrations and conditions may vary. Information and photo are courtesy of the Tissue Profiling group, SciLifeLab Stockholm. Neutralization: To yield one-half maximal inhibition [ND₅₀] of the biological activity of Murine IL-6 (3 pg/ml), a concentration of 0.24-0.36 µg/ml of this antibody is required. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.