

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Anti-Murine IP-10 Antibody, Rabbit, Polyclonal ABT-ABG10316-U050

Artikelname	Anti-Murine IP-10 Antibody, Rabbit, Polyclonal
Artikelnummer	ABT-ABG10316-U050
Hersteller Artikelnummer	ABG10316-U050
Alternativnummer	ABT-ABG10316-U050-50UG
Hersteller	Abcepta
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA, IHC, WB
Spezies Reaktivität	Mouse
Klonalität	Polyclonal
Reinheit	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant mIP-10. Anti-Murine IP-10 specific antibody was purified by affinity chromatography employing immobilized mIP-10 matrix.
Formulierung	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Anwendungsbeschreibung

WesternBlot: To detect mIP-10 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mIP-10 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect mIP-10 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Murine IP-10 (61-081BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mIP-10.. Immunohistochemistry: This antibody stained colchicine injected mouse brain (including the hippocampus region) tissue. The primary antibody was incubated at 1.0 mg/ml overnight at 4°C. This was followed by a peroxidase conjugated secondary antibody and then a fluorescein Tyramide Signal Amplification (TSA(TM)) reagent. Optimal concentrations and conditions may vary. . Neutralization: To yield one-half maximal inhibition [ND<span style=font-size: 16px,>50] of the biological activity of mIP-10 (100.0 ng/ml), a concentration of 10.0 µg/ml of this antibody is required.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.