

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Murine Leptin Antibody, Rabbit, Polyclonal ABT-ABG10340-U100

Artikelname	Anti-Murine Leptin Antibody, Rabbit, Polyclonal
Artikelnummer	ABT-ABG10340-U100
Hersteller Artikelnummer	ABG10340-U100
Alternativnummer	ABT-ABG10340-U100-100UG
Hersteller	Abcepta
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA, IHC, WB
Spezies Reaktivität	Mouse
Klonalität	Polyclonal
Reinheit	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant mLeptin. Anti-Murine Leptin specific antibody was purified by affinity chromatography employing immobilized mLeptin matrix.
Formulierung	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Anwendungsbeschreibung

WesternBlot: To detect mLeptin by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mLeptin is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect mLeptin by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Murine Leptin (61-085BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mLeptin.. Immunohistochemistry: This antibody stained 4% PFA, paraffin-embedded sections of *Trichuris muris* infected murine cecum and large intestine tissue. The primary antibody was incubated at 0.20 µg/mL overnight at 4C and the secondary antibody was a cyanine-3 conjugated Donkey anti-Rabbit (Jackson ImmunoResearch). Heat induced antigen retrieval with a 100mM citric acid was used. Information and photo are courtesy of David Artis, University of Pennsylvania. Optimal concentrations and conditions may vary.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.