

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Human LIGHT Antibody, Rabbit, Polyclonal ABT-ABG10346-U100

Artikelname	Anti-Human LIGHT Antibody, Rabbit, Polyclonal
Artikelnummer	ABT-ABG10346-U100
Hersteller Artikelnummer	ABG10346-U100
Alternativnummer	ABT-ABG10346-U100-100UG
Hersteller	Abcepta
Wirt	Rabbit
Kategorie	Antikörper
Applikation	ELISA, IHC, WB
Spezies Reaktivität	Human
Klonalität	Polyclonal
Reinheit	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hLIGHT . Anti-Human LIGHT specific antibody was purified by affinity chromatography employing immobilized hLIGHT matrix.
Formulierung	A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.
Antibody Type	Polyclonal Antibody

Anwendungsbeschreibung	<p>WesternBlot: To detect hLIGHT by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hLIGHT is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hLIGHT by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems Biotinylated Anti-Human LIGHT (60-206BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hLIGHT..</p> <p>Immunohistochemistry: This antibody stained formalin-fixed, paraffin-embedded sections of human normal placenta. The recommended concentration is 2.5 µg/mL with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 8.0 buffer is recommended. Optimal concentrations and conditions may vary.

 Tissue samples were provided by the Cooperative Human Tissue Network, which is funded by the National Cancer Institute.
 Information and photo are courtesy of Flagship Biosciences, LLC - Histology Services. .</p> <p>Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.</p>
------------------------	---