

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Biotinylated Anti-Human MIA Antibody, Rabbit, Polyclonal ABT-ABG10388-U025**

|                          |   |
|--------------------------|---|
| Artikelname              | Biotinylated Anti-Human MIA Antibody, Rabbit, Polyclonal  |
| Artikelnummer            | ABT-ABG10388-U025   |
| Hersteller Artikelnummer | ABG10388-U025   |
| Alternativnummer         | ABT-ABG10388-U025-25UG  |
| Hersteller               | Abcepta   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | ELISA, WB   |
| Spezies Reaktivität      | Human   |
| Klonalität               | Polyclonal  |
| Reinheit                 | Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hMIA. Anti-Human MIA specific antibody was purified by affinity chromatography and then biotinylated. |
| Formulierung             | A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.  |
| Antibody Type            | Polyclonal Antibody   |

|                        |  |
|------------------------|--|
| Anwendungsbeschreibung | <p>WesternBlot: To detect hMIA by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMIA is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.. Sandwich: To detect hMIA by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.25 - 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems Polyclonal Anti-Human MIA (60-219P) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMIA.. Direct: To detect hMIA by direct ELISA (using 100 µl/well antibody solution) a concentration of 0.25 - 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMIA.. Reconstitution: Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.</p> |
|------------------------|--|