

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Recombinant CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) Antibody, IgG1, Clone: [rCFTR/1342], Mouse, Monoclonal NBT-1080-MSM9-P1ABX

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) Antibody, IgG1, Clone: [rCFTR/1342], Mouse, Monoclonal   |
| Artikelnummer            | NBT-1080-MSM9-P1ABX   |
| Hersteller Artikelnummer | 1080-MSM9-P1ABX   |
| Alternativnummer         | NBT-1080-MSM9-P1ABX-100   |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human, Mouse  |
| Immunogen                | Recombinant full-length human CFTR protein.   |
| Produktbeschreibung      | Recognizes a protein of 165-170kDa, identified as cystic fibrosis transmembrane conductance regulator (CFTR). CFTR is composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structurally similar t... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [rCFTR/1342]  |
| Molekulargewicht         | 165-170kDa  |
| Isotyp                   | IgG1  |

|                        |   |
|------------------------|---|
| NCBI                   | <a href="#">1080</a>  |
| UniProt                | <a href="#">P13569</a>  |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Recombinant Monoclonal Antibody   |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95degC followed by cooling at RT for 20 minutes)Optimal dilu |