

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### HCG-beta Antibody, IgG1, Clone: [SPM529], Mouse, Monoclonal NBT-1082-MSM2X-P1ABX

|                          |   |
|--------------------------|---|
| Artikelname              | HCG-beta Antibody, IgG1, Clone: [SPM529], Mouse, Monoclonal   |
| Artikelnummer            | NBT-1082-MSM2X-P1ABX  |
| Hersteller Artikelnummer | 1082-MSM2X-P1ABX  |
| Alternativnummer         | NBT-1082-MSM2X-P1ABX-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant hCG beta protein  |
| Produktbeschreibung      | This MAb reacts with a protein of 22kDa, identified as beta sub-unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in ... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [SPM529]  |
| Molekulargewicht         | 22kDa   |
| Isotyp                   | IgG1  |
| NCBI                     | <a href="#">1082</a>  |

|                        |  |
|------------------------|--|
| UniProt                | <a href="#">P01233</a>   |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Monoclonal Antibody  |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml, 30 min RT)<br>(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution for a spec |