

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Recombinant Carboxypeptidase A1 / CPA1 (Pancreatic Cancer Marker) Antibody, IgG1, Clone: [rCPA1/8688], Mouse, Monoclonal NBT-1357-MSM14-P1ABX**

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant Carboxypeptidase A1 / CPA1 (Pancreatic Cancer Marker) Antibody, IgG1, Clone: [rCPA1/8688], Mouse, Monoclonal  |
| Artikelnummer            | NBT-1357-MSM14-P1ABX  |
| Hersteller Artikelnummer | 1357-MSM14-P1ABX  |
| Alternativnummer         | NBT-1357-MSM14-P1ABX-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human CPA1 protein  |
| Produktbeschreibung      | Human pancreatic procarboxypeptidase A exists as three different active forms, two of which are designated carboxypeptidase A1 (CPA1) and carboxypeptidase A2 (CPA2). CPA1, also known as CPA, is a 419 amino acid secreted monomeric protein that is high... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [rCPA1/8688]  |
| Molekulargewicht         | 47kDa   |
| Isotyp                   | IgG1  |

|                        |  |
|------------------------|--|
| NCBI                   | <a href="#">1357</a>   |
| UniProt                | <a href="#">P15085</a>   |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Recombinant Monoclonal Antibody  |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml),,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 |