

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Recombinant Estrogen Receptor, alpha (Marker of Estrogen Dependence) Antibody, Clone: [ESR1/7444R], Rabbit, Monoclonal NBT-2099-RBM44-P0**

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant Estrogen Receptor, alpha (Marker of Estrogen Dependence) Antibody, Clone: [ESR1/7444R], Rabbit, Monoclonal  |
| Artikelnummer            | NBT-2099-RBM44-P0   |
| Hersteller Artikelnummer | 2099-RBM44-P0   |
| Alternativnummer         | NBT-2099-RBM44-P0-20,NBT-2099-RBM44-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant fragment (around aa495-595) of human ER alpha protein (exact sequence is proprietary)   |
| Produktbeschreibung      | Estrogen receptor (ER) belongs to the steroid receptor superfamily of nuclear receptors. It is a protein with 553 amino acids. The receptor molecule has three domains, i.e. the central DNA-binding domain, the hormone-binding domain at the C-terminal,... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [ESR1/7444R]  |
| Molekulargewicht         | ~67kDa  |

|                        |  |
|------------------------|--|
| NCBI                   | <a href="#">2099</a>   |
| UniProt                | <a href="#">P03372</a>   |
| Formulierung           | 200ug/ml of Ab purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Recombinant Monoclonal Antibody  |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes),Optimal dilutio |