

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### EpCAM / CD326 (Epithelial Marker) Antibody, Rabbit, Polyclonal NBT-4072-RBP17-P0

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | EpCAM / CD326 (Epithelial Marker) Antibody, Rabbit, Polyclonal                                                                                                                                                                                                |
| Artikelnummer            | NBT-4072-RBP17-P0                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | 4072-RBP17-P0                                                                                                                                                                                                                                                 |
| Alternativnummer         | NBT-4072-RBP17-P0-20,NBT-4072-RBP17-P0-100                                                                                                                                                                                                                    |
| Hersteller               | NeoBiotechnologies                                                                                                                                                                                                                                            |
| Wirt                     | Rabbit                                                                                                                                                                                                                                                        |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | FC, IF, WB                                                                                                                                                                                                                                                    |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant human TACSTD1 protein                                                                                                                                                                                                                             |
| Produktbeschreibung      | EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (EpCAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majori... |
| Klonalität               | Polyclonal                                                                                                                                                                                                                                                    |
| Molekulargewicht         | 40-43kDa                                                                                                                                                                                                                                                      |
| NCBI                     | <a href="#">4072</a>                                                                                                                                                                                                                                          |
| UniProt                  | <a href="#">P16422</a>                                                                                                                                                                                                                                        |

|                        |                                                                                                                                                                                                                                                               |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Formulierung           | 200ug/ml of Ab purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.                                                                                                                      |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), ,Western Blotting (1-2ug/ml), ,Immunohistology (Formalin-fixed) (1-2ug/ml for 30 min at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate Buffer, |