

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Recombinant Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody, Clone: [BCL2/2210R], Rabbit, Monoclonal NBT-596-RBM10-P0

Artikelname	Recombinant Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody, Clone: [BCL2/2210R], Rabbit, Monoclonal
Artikelnummer	NBT-596-RBM10-P0
Hersteller Artikelnummer	596-RBM10-P0
Alternativnummer	NBT-596-RBM10-P0-20,NBT-596-RBM10-P0-100
Hersteller	NeoBiotechnologies
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human bcl-2 protein
Produktbeschreibung	This antibody recognizes a protein of 25-26kDa, identified as the bcl-2 lpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of bcl-2 lpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular ly...
Klonalität	Monoclonal
Klon-Bezeichnung	[BCL2/2210R]
Molekulargewicht	25-26kDa
NCBI	<a href="#">596</a>

UniProt	<a href="#">P10415</a>
Formulierung	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody Type	Recombinant Monoclonal Antibody
Anwendungsbeschreibung	Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 7.5-8.5, for 45 min at 95&degC followed by cooling at RT for 20 minutes)Opt