

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TIM3 / HAVCR2 / CD366 (Effector T-Cell Marker) Antibody, IgG2b, Clone: [TIM3/4030], Mouse, Monoclonal NBT-84868-MSM10-P0

| | |
|--------------------------|---|
| Artikelname | TIM3 / HAVCR2 / CD366 (Effector T-Cell Marker) Antibody, IgG2b, Clone: [TIM3/4030], Mouse, Monoclonal |
| Artikelnummer | NBT-84868-MSM10-P0 |
| Hersteller Artikelnummer | 84868-MSM10-P0 |
| Alternativnummer | NBT-84868-MSM10-P0-20,NBT-84868-MSM10-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of human TIM3 protein (around aa 22-202) (exact sequence is proprietary) |
| Produktbeschreibung | TIMs are type I transmembrane glycoproteins with one Ig-like V-type domain and a Ser/Thr-rich mucin stalk. TIM-3 is expressed on the surface of effector T cel tonsil, lymph node or spleen. Is (CD4+Th1 and CD8+Tc1) but not on helper T cells (CD4+Th2 a... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [TIM3/4030] |
| Molekulargewicht | 60kDa |

| | |
|------------------------|---|
| Isotyp | IgG2b |
| NCBI | 84868 |
| UniProt | Q8TDQ0 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |