

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Proliferation Marker Antibody, IgG2a, Clone: [JC1], Mouse, Monoclonal NBT-MSM1-3329-P1ABX

| | |
|--------------------------|---|
| Artikelname | Proliferation Marker Antibody, IgG2a, Clone: [JC1], Mouse, Monoclonal |
| Artikelnummer | NBT-MSM1-3329-P1ABX |
| Hersteller Artikelnummer | MSM1-3329-P1ABX |
| Alternativnummer | NBT-MSM1-3329-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Pokeweed-stimulated lymphocytes |
| Produktbeschreibung | JC1 detects a nuclear antigen, distinct from Ki67, present in proliferating cells. It is useful as a marker of proliferation in cases such as squamous cell carcinoma where the Ki67 index cannot be determined.... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [JC1] |
| Molekulargewicht | 212kDa and 123kDa |
| Isotyp | IgG2a |

| | |
|------------------------|--|
| NCBI | Not Known |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followe |