

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Beta-2 Microglobulin Antibody [B2M/961], Unconjugated, Mouse, Monoclonal PRS-33-458

| | |
|--------------------------|--|
| Artikelname | Beta-2 Microglobulin Antibody [B2M/961], Unconjugated, Mouse, Monoclonal |
| Artikelnummer | PRS-33-458 |
| Hersteller Artikelnummer | 33-458 |
| Alternativnummer | PRS-33-458-100 |
| Hersteller | ProSci |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC-P |
| Spezies Reaktivität | Human, Primate |
| Immunogen | Full length recombinant human protein was used as the immunogen for the Beta-2 Microglobulin antibody. |
| Konjugation | Unconjugated |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [B2M/961] |
| NCBI | 567 |
| UniProt | P61769 |

| | |
|------------------------|--|
| Puffer | PBS with 0.1 mg/ml BSA and 0.05% sodium azide |
| Formulierung | Liquid |
| Application Verdünnung | Optimal dilutions for each application to be determined by the researcher |
| Anwendungsbeschreibung | <p>Flow Cytometry: 0.5-1 ug/million cells in 0.1ml Immunofluorescence: 0.5-1 ug/ml Immunohistochemistry (FFPE): 0.5-1 ug/ml for 30 min at RT (1) Prediluted format: incubate for 30 min at RT (2) Optimal dilution of the Beta-2 Microglobulin antibody should be determined by the researcher.</p> <p>1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.</p> <p>2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.</p> |