

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TIGIT Antibody [4A10], IgG1, Unconjugated, Mouse, Monoclonal PRS-RF16054

| | |
|--------------------------|---|
| Artikelname | TIGIT Antibody [4A10], IgG1, Unconjugated, Mouse, Monoclonal |
| Artikelnummer | PRS-RF16054 |
| Hersteller Artikelnummer | RF16054 |
| Alternativnummer | PRS-RF16054-0.02, PRS-RF16054-0.1 |
| Hersteller | ProSci |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, FC, ICC, IF, IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | TIGIT antibody was raised against the extracellular domain of human TIGIT |
| Konjugation | Unconjugated |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [4A10] |
| Molekulargewicht | Predicted: 26 kDa Observed: 47 kDa |
| Isotyp | IgG1 |
| NCBI | 201633 |

| | |
|------------------------|---|
| UniProt | Q495A1 |
| Puffer | TIGIT Antibody is supplied in PBS containing 0.02% sodium azide and 50% glycerol. |
| Formulierung | Liquid |
| Application Verdünnung | Optimal dilutions for each application to be determined by the researcher. |
| Anwendungsbeschreibung | TIGIT antibody can be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 1 µg/mL. For flow cytometry at 1 µg/ml. For immunocytochemistry at 1 µg/mL. For Western blot at 1 µg/mL. Antibody validated: Western Blot in human samples, Immunohistochemistry in human samples, Immunocytochemistry in human samples, Immunofluorescence in human samples and Flow Cytometry in human samples. All other applications and species not yet tested. |