

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Exodus-2 Antibody, Unconjugated, Rabbit, Polyclonal PRS-XP-5128

|                          |   |
|--------------------------|---|
| Artikelname              | Exodus-2 Antibody, Unconjugated, Rabbit, Polyclonal   |
| Artikelnummer            | PRS-XP-5128   |
| Hersteller Artikelnummer | XP-5128   |
| Alternativnummer         | PRS-XP-5128-0.1   |
| Hersteller               | ProSci  |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | ELISA, NeA, WB  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hExodus-2 (human Exodus-2). |
| Konjugation              | Unconjugated  |
| Klonalität               | Polyclonal  |
| Konzentration            | batch dependent   |
| NCBI                     | <a href="#">6366</a>  |
| UniProt                  | <a href="#">000585</a>  |
| Formulierung             | Lyophilized   |
| Application Verdünnung   | Centrifuge vial prior to opening.   |

Anwendungsbeschreibung

Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of hExodus-2 (100.00 ng/mL), a concentration of 6.0 - 8.0 µg/mL of this antibody is required. ELISA: To detect hExodus-2 by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hExodus 2. Sandwich: To detect hExodus-2 by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Human Exodus-2 (XP-5128Bt) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hExodus-2. Western Blot: To detect hExodus-2 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hExodus-2 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.