

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 18(DE-K18), CF405S conjugate, 0.1mg/mL, Clone: [DE-K18], Mouse, Monoclonal BOT-BNC040563-500

Artikelname	Cytokeratin 18(DE-K18), CF405S conjugate, 0.1mg/mL, Clone: [DE-K18], Mouse, Monoclonal
Artikelnummer	BOT-BNC040563-500
Hersteller Artikelnummer	BNC040563-500
Alternativnummer	BOT-BNC040563-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Canine, Feline, Human
Immunogen	Cytoskeletal preparation extracted from the human vulvar squamous carcinoma cell line A431
Konjugation	CF405S
Produktbeschreibung	This MAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs t...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[DE-K18]
Molekulargewicht	45 kDa
UniProt	P05783
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with rat, others not known Immunohistology formalin-fixed 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user