

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Histone H1(1415-1), CF405S conjugate, 0.1mg/mL, Clone: [1415-1], Mouse, Monoclonal BOT-BNC040580-100**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Histone H1(1415-1), CF405S conjugate, 0.1mg/mL, Clone: [1415-1], Mouse, Monoclonal                                                                                                                                                                            |
| Artikelnummer            | BOT-BNC040580-100                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | BNC040580-100                                                                                                                                                                                                                                                 |
| Alternativnummer         | BOT-BNC040580-100-100UL                                                                                                                                                                                                                                       |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | FC, IF, IHC, WB                                                                                                                                                                                                                                               |
| Spezies Reaktivität      | Human, Mouse, Rat                                                                                                                                                                                                                                             |
| Immunogen                | Nuclei of human leukemia biopsy cells                                                                                                                                                                                                                         |
| Konjugation              | CF405S                                                                                                                                                                                                                                                        |
| Produktbeschreibung      | Eukaryotic histones are basic, water-soluble nuclear proteins that form hetero-octameric nucleosome particles. They wrap 146 base pairs of DNA in a left-handed supealpha-helicalturn sequentially to form chromosomal fiber. Two molecules of each of the... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [1415-1]                                                                                                                                                                                                                                                      |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | ~30 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| UniProt                | Multiple                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |