

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 8(TS1), CF405S conjugate, 0.1mg/mL, Clone: [TS1], Mouse, Monoclonal BOT-BNC040627-500**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Cytokeratin 8(TS1), CF405S conjugate, 0.1mg/mL, Clone: [TS1], Mouse, Monoclonal                                                                                                                                                                               |
| Artikelnummer            | BOT-BNC040627-500                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | BNC040627-500                                                                                                                                                                                                                                                 |
| Alternativnummer         | BOT-BNC040627-500-500UL                                                                                                                                                                                                                                       |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | FC, IF, IHC, WB                                                                                                                                                                                                                                               |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Keratin preparation from a human carcinoma                                                                                                                                                                                                                    |
| Konjugation              | CF405S                                                                                                                                                                                                                                                        |
| Produktbeschreibung      | Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins, and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adeno... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [TS1]                                                                                                                                                                                                                                                         |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | 52.5 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| UniProt                | <a href="#">P05787</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with rat or ferret, others not known Immunohistology formalin-fixed 0.5-1.0 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |