

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 19(BA17), CF405S conjugate, 0.1mg/mL, Clone: [BA17], Mouse, Monoclonal BOT-BNC040666-100

Artikelname	Cytokeratin 19(BA17), CF405S conjugate, 0.1mg/mL, Clone: [BA17], Mouse, Monoclonal
Artikelnummer	BOT-BNC040666-100
Hersteller Artikelnummer	BNC040666-100
Alternativnummer	BOT-BNC040666-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC, WB
Spezies Reaktivität	Human, Mouse
Immunogen	Human mammary epithelial organoids.
Konjugation	CF405S
Produktbeschreibung	This Ab reacts with the rod domain of human cytokeratin 19 (CK19), a polypeptide of 40 kDa. CK19 is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus,...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[BA17]
Molekulargewicht	40 kDa
UniProt	P08727
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user