

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **PCNA(PCNA/694), CF405S conjugate, 0.1mg/mL, Clone: [PCNA/694], Mouse, Monoclonal BOT-BNC040694-100**

|                          |   |
|--------------------------|---|
| Artikelname              | PCNA(PCNA/694), CF405S conjugate, 0.1mg/mL, Clone: [PCNA/694],<br>Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNC040694-100   |
| Hersteller Artikelnummer | BNC040694-100   |
| Alternativnummer         | BOT-BNC040694-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full length human PCNA protein  |
| Konjugation              | CF405S  |
| Produktbeschreibung      | Recognizes a non-histone protein of 36 kDa, which is identified as proliferating cell nuclear antigen (PCNA). It is also known as cyclin or polymerase delta auxiliary protein. Elevated expression of PCNA/cyclin has been shown in the nucleus during la... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [PCNA/694]  |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 36 kDa   |
| UniProt                | <a href="#">P12004</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Predicted to show broad species reactivity Optimal dilution for a specific application should be determined by user</p> |