

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD66 (CEA)(COL-1 + CEA31 + C66/261), CF405S conjugate, 0.1mg/mL, Clone: [COL-1 CEA31 C66/261], Mouse, Monoclonal BOT-BNC040747-500

| | |
|--------------------------|---|
| Artikelname | CD66 (CEA)(COL-1 + CEA31 + C66/261), CF405S conjugate, 0.1mg/mL, Clone: [COL-1 CEA31 C66/261], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040747-500 |
| Hersteller Artikelnummer | BNC040747-500 |
| Alternativnummer | BOT-BNC040747-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Human colon carcinoma extract (COL-1 & CEA31), Recombinant full-length human CEA protein (C66/261) |
| Konjugation | CF405S |
| Produktbeschreibung | This antibody recognizes proteins of 80-200 kDa, identified as different members of CEA family. CEA is synthesized during development in the fetal gut and is re-expressed in increased amounts in intestinal carcinomas and several other tumors. This MA... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [COL-1 CEA31 C66/261] |
| Molekulargewicht | 80-200 kDa |
| UniProt | P06731 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |