

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD8(C8/468 + C8/144B), CF405S conjugate, 0.1mg/mL, Clone: [C8/468 C8/144B], Mouse, Monoclonal BOT-BNC040750-100**

Artikelname	CD8(C8/468 + C8/144B), CF405S conjugate, 0.1mg/mL, Clone: [C8/468 C8/144B], Mouse, Monoclonal
Artikelnummer	BOT-BNC040750-100
Hersteller Artikelnummer	BNC040750-100
Alternativnummer	BOT-BNC040750-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Human CD8 recombinant protein (C8/468), A 13 amino acid synthetic peptide from the C-terminal cytoplasmic domain of alpha chain of human CD8 molecule (C8/144B)
Konjugation	CF405S
Produktbeschreibung	CD8 is a cell surface receptor expressed either as a heterodimer with the CD8 beta chain (CD8 alpha/beta) or as a homodimer (CD8 alpha/alpha). A majority of thymocytes and a subpopulation of mature T cells and NK cells express CD8a. CD8 binds to MHC ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[C8/468 C8/144B]
Molekulargewicht	32 kDa
UniProt	<a href="#">P01732</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Not recommended for flow cytometry Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user