

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human IgG Immunoglobulin(ICO-97), CF405S conjugate, 0.1mg/mL, Clone: [ICO-97], Mouse, Monoclonal BOT-BNC040758-100

| | |
|--------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Human IgG Immunoglobulin(ICO-97), CF405S conjugate, 0.1mg/mL, Clone: [ICO-97], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC040758-100 |
| Hersteller Artikelnummer | BNC040758-100 |
| Alternativnummer | BOT-BNC040758-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Purified human Ig Gamma Chain |
| Konjugation | CF405S |
| Produktbeschreibung | Recognizes a protein of 75 kDa, identified as i heavy chain of human immunoglobulins. It does not cross-react with i (IgA), i (IgM), i μ (IgE), or i' (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. This MAb is useful in the ide... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [ICO-97] |
| Molekulargewicht | 75 kDa |

| | |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt | P01857 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |