

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD20(IGEL/773), CF405S conjugate, 0.1mg/mL, Clone: [IGEL/773], Mouse, Monoclonal BOT-BNC040773-500**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | CD20(IGEL/773), CF405S conjugate, 0.1mg/mL, Clone: [IGEL/773], Mouse, Monoclonal                                                                                                                                                                              |
| Artikelnummer            | BOT-BNC040773-500                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | BNC040773-500                                                                                                                                                                                                                                                 |
| Alternativnummer         | BOT-BNC040773-500-500UL                                                                                                                                                                                                                                       |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | FC, IF, IHC                                                                                                                                                                                                                                                   |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant human MS4A1 protein                                                                                                                                                                                                                               |
| Konjugation              | CF405S                                                                                                                                                                                                                                                        |
| Produktbeschreibung      | Recognizes a protein of 30-33 kDa, which is identified as CD20. It is a non-Ig differentiation antigen of B-cells and its expression is restricted to normal and neoplastic B-cells, being absent from all other leukocytes and tissues. CD20 is expressed... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [IGEL/773]                                                                                                                                                                                                                                                    |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | 33-37 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| UniProt                | <a href="#">P11836</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |