

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Ep-CAM / CD326 (Epithelial Marker) (EGP40/1556R), CF405S conjugate, 0.1mg/mL, Clone: [EGP40/1556R], Rabbit, Monoclonal BOT-BNC041556-100**

Artikelname	Ep-CAM / CD326 (Epithelial Marker) (EGP40/1556R), CF405S conjugate, 0.1mg/mL, Clone: [EGP40/1556R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC041556-100
Hersteller Artikelnummer	BNC041556-100
Alternativnummer	BOT-BNC041556-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human TACSTD1 protein
Konjugation	CF405S
Produktbeschreibung	EGP40 is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majo...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[EGP40/1556R]

Molekulargewicht	40-43 kDa
UniProt	<a href="#">P16422</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user