

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin, Basic (Type II or HMW) (Epithelial Marker) (KRTH/1576R), CF405S conjugate, 0.1mg/mL, Clone: [KRTH/1576R], Rabbit, Monoclonal BOT-BNC041576-100**

|                          |   |
|--------------------------|---|
| Artikelname              | Cytokeratin, Basic (Type II or HMW) (Epithelial Marker) (KRTH/1576R), CF405S conjugate, 0.1mg/mL, Clone: [KRTH/1576R], Rabbit, Monoclonal   |
| Artikelnummer            | BOT-BNC041576-100   |
| Hersteller Artikelnummer | BNC041576-100   |
| Alternativnummer         | BOT-BNC041576-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human KRT76 protein   |
| Konjugation              | CF405S  |
| Produktbeschreibung      | This MAb recognizes basic (Type II or HMW) cytokeratins, which include 67 kDa (CK1), 64 kDa (CK3), 59 kDa (CK4), 58 kDa (CK5), 56 kDa (CK6), 52 kDa (CK8). Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |

|                        |  |
|------------------------|--|
| Klon-Bezeichnung       | [KRTH/1576R]   |
| Molekulargewicht       | 52-67 kDa  |
| UniProt                | <a href="#">Q01546</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.25-0.5 ug/mL Immunofluorescence 1-2 ug/mL Western blotting 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |