

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Spectrin beta III (SPTBN2) (SPTBN2/1583), CF405S conjugate, 0.1mg/mL, Clone: [SPTBN2/1583], Mouse, Monoclonal BOT-BNC041583-500

| | |
|--------------------------|--|
| Artikelname | Spectrin beta III (SPTBN2) (SPTBN2/1583), CF405S conjugate, 0.1mg/mL, Clone: [SPTBN2/1583], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC041583-500 |
| Hersteller Artikelnummer | BNC041583-500 |
| Alternativnummer | BOT-BNC041583-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human SPTBN2 fragment (aa356-475) (exact sequence is proprietary) |
| Konjugation | CF405S |
| Produktbeschreibung | Spectrin is an actin binding protein that is a major component of the plasma membrane skeleton. Spectrins function as membrane organizers and stabilizers by forming dimers, tetramers and higher polymers. Vertebrate spectrins have two alpha-subunits (...) |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [SPTBN2/1583] |
| Molekulargewicht | 246 kDa |
| UniProt | O15020 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |