

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

OCT-2 (POU2F2) (B-Cell Marker) (OCT2/2136), CF405S conjugate, 0.1mg/mL, Clone: [OCT2/2136], Mouse, Monoclonal BOT-BNC042136-500

Artikelname	OCT-2 (POU2F2) (B-Cell Marker) (OCT2/2136), CF405S conjugate, 0.1mg/mL, Clone: [OCT2/2136], Mouse, Monoclonal
Artikelnummer	BOT-BNC042136-500
Hersteller Artikelnummer	BNC042136-500
Alternativnummer	BOT-BNC042136-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant fragment of human OCT2 protein (around aa 112-297) (exact sequence is proprietary)
Konjugation	CF405S
Produktbeschreibung	Oct-2 is a transcription factor of the POU homeo-domain family that binds to the Ig gene octamer sites, regulating B-cell-specific genes. Oct-2 expression can be used as a marker of B-cell lineage and differentiation. Germinal center B-cells, mantle ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[OCT2/2136]
Molekulargewicht	60 kDa
UniProt	P09086
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user