

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Bcl-2 (Apoptosis & Follicular Lymphoma Marker) (rBCL2/782), CF405S conjugate, 0.1mg/mL, Clone: [rBCL2/782], Mouse, Monoclonal BOT-BNC042221-500**

|                          |   |
|--------------------------|---|
| Artikelname              | Bcl-2 (Apoptosis & Follicular Lymphoma Marker) (rBCL2/782), CF405S conjugate, 0.1mg/mL, Clone: [rBCL2/782], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNC042221-500   |
| Hersteller Artikelnummer | BNC042221-500   |
| Alternativnummer         | BOT-BNC042221-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human BCL2 protein  |
| Konjugation              | CF405S  |
| Produktbeschreibung      | This antibody recognizes a protein of 25-26 kDa, identified as Bcl-2. This antibody shows no cross-reaction with Bcl-x or Bax proteins. Bcl-2 suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural ... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |

|                        |  |
|------------------------|--|
| Klon-Bezeichnung       | [rBCL2/782]  |
| Molekulargewicht       | 25-26 kDa  |
| UniProt                | <a href="#">P10415</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 1 mM EDTA Buffer pH 8.0 for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |