

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

**Cytokeratin 19 (KRT19) (Pancreatic Stem Cell Marker) (rKRT19/799),  
CF405S conjugate, 0.1mg/mL, IgG1, Clone: [rKRT19/799], Mouse,  
Monoclonal  
BOT-BNC042304-100**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Cytokeratin 19 (KRT19) (Pancreatic Stem Cell Marker) (rKRT19/799), CF405S conjugate, 0.1mg/mL, IgG1, Clone: [rKRT19/799], Mouse, Monoclonal                                                                                                                   |
| Artikelnummer            | BOT-BNC042304-100                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | BNC042304-100                                                                                                                                                                                                                                                 |
| Alternativnummer         | BOT-BNC042304-100-100UL                                                                                                                                                                                                                                       |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | FC, IHC, WB                                                                                                                                                                                                                                                   |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant full-length human KRT19 protein                                                                                                                                                                                                                   |
| Konjugation              | CF405S                                                                                                                                                                                                                                                        |
| Produktbeschreibung      | This MAb reacts with the rod domain of human cytokeratin-19 (CK19), a polypeptide of 40 kDa. Its epitope maps between amino acid 312-335. CK19 is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, ... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.1 mg/mL                                                                                                                                                                                                                                                     |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Klon-Bezeichnung       | [rKRT19/799]                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Molekulargewicht       | 40 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Isotyp                 | IgG1                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| UniProt                | <a href="#">P08727</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with rat or ferret Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 1-2 ug/mL Optimal dilution for a specific application should be determined by user</p> |