

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 6A (KRT6A) (Basal Cell Marker) (KRT6A/2368), CF405S conjugate, 0.1mg/mL, Clone: [KRT6A/2368], Mouse, Monoclonal BOT-BNC042368-100**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Cytokeratin 6A (KRT6A) (Basal Cell Marker) (KRT6A/2368), CF405S conjugate, 0.1mg/mL, Clone: [KRT6A/2368], Mouse, Monoclonal                                                                                                                                   |
| Artikelnummer            | BOT-BNC042368-100                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | BNC042368-100                                                                                                                                                                                                                                                 |
| Alternativnummer         | BOT-BNC042368-100-100UL                                                                                                                                                                                                                                       |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | FC, IHC                                                                                                                                                                                                                                                       |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant full-length human Cytokeratin 6A (KRT6A) protein                                                                                                                                                                                                  |
| Konjugation              | CF405S                                                                                                                                                                                                                                                        |
| Produktbeschreibung      | This MAb recognizes a protein of 56 kDa, identified as cytokeratin 6A (KRT6A). In humans, multiple isoforms of Cytokeratin 6 (6A-6F), encoded by several highly homologous genes, have distinct tissue expression patterns. Cytokeratin 6A is the dominant... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [KRT6A/2368]                                                                                                                                                                                                                                                  |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | 56 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| UniProt                | <a href="#">P02538</a>                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |