

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **GP2 (Glycoprotein 2) / ZAP75 (GP2/2569R), CF405S conjugate, 0.1mg/mL, Clone: [GP2/2569R], Rabbit, Monoclonal BOT-BNC042569-500**

Artikelname	GP2 (Glycoprotein 2) / ZAP75 (GP2/2569R), CF405S conjugate, 0.1mg/mL, Clone: [GP2/2569R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC042569-500
Hersteller Artikelnummer	BNC042569-500
Alternativnummer	BOT-BNC042569-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human GP2 protein fragment (around aa35-179) (exact sequence is proprietary)
Konjugation	CF405S
Produktbeschreibung	GP2 (glycoprotein 2), also known as ZAP75, is a 537 amino acid secreted protein. It is an integral membrane protein that is secreted from intracellular zymogen granules and associates with the plasma membrane via glycosylphosphatidylinositol (GPI) li...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[GP2/2569R]
Molekulargewicht	59 kDa
UniProt	<a href="#">P55259</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with mouse ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user