

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 18 (KRT18) (KRT18/2808R), CF405S conjugate, 0.1mg/mL, Clone: [KRT18/2808R], Rabbit, Monoclonal BOT-BNC042808-100**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Cytokeratin 18 (KRT18) (KRT18/2808R), CF405S conjugate, 0.1mg/mL, Clone: [KRT18/2808R], Rabbit, Monoclonal                                                                                                                                                    |
| Artikelnummer            | BOT-BNC042808-100                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | BNC042808-100                                                                                                                                                                                                                                                 |
| Alternativnummer         | BOT-BNC042808-100-100UL                                                                                                                                                                                                                                       |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Rabbit                                                                                                                                                                                                                                                        |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | FC, IF, IHC, WB                                                                                                                                                                                                                                               |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant full-length human KRT18 protein                                                                                                                                                                                                                   |
| Konjugation              | CF405S                                                                                                                                                                                                                                                        |
| Produktbeschreibung      | This MAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs t... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [KRT18/2808R]                                                                                                                                                                                                                                                 |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | 45 kDa                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| UniProt                | <a href="#">P05783</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Western Blot 0.5-2 ug/mL Optimal dilution for a specific application should be determined by user</p> |