

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Golgi Complex (Marker for Human Cells) (GLG1/2829R), CF405S conjugate, 0.1mg/mL, Clone: [GLG1/2829R], Rabbit, Monoclonal BOT-BNC042829-100**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Golgi Complex (Marker for Human Cells) (GLG1/2829R), CF405S conjugate, 0.1mg/mL, Clone: [GLG1/2829R], Rabbit, Monoclonal                                                                                                                                      |
| Artikelnummer            | BOT-BNC042829-100                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | BNC042829-100                                                                                                                                                                                                                                                 |
| Alternativnummer         | BOT-BNC042829-100-100UL                                                                                                                                                                                                                                       |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Rabbit                                                                                                                                                                                                                                                        |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC                                                                                                                                                                                                                                                           |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | SU-DHL-1 large cell lymphoma cells.                                                                                                                                                                                                                           |
| Konjugation              | CF405S                                                                                                                                                                                                                                                        |
| Produktbeschreibung      | This MAb recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining ... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [GLG1/2829R]                                                                                                                                                                                                                                                  |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | Not Known                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| UniProt                | Not Known                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with mouse or rat Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user</p> |