

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cyclin B1 (G2- & M-phase Cyclin)(CCNB1/2978R), CF405S conjugate, 0.1mg/mL, Clone: [CCNB1/2978R], Rabbit, Monoclonal BOT-BNC042978-100**

Artikelname	Cyclin B1 (G2- & M-phase Cyclin)(CCNB1/2978R), CF405S conjugate, 0.1mg/mL, Clone: [CCNB1/2978R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC042978-100
Hersteller Artikelnummer	BNC042978-100
Alternativnummer	BOT-BNC042978-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Spezies Reaktivität	Hamster, Human, Mouse
Immunogen	Recombinant full-length human CCNB1 protein
Konjugation	CF405S
Produktbeschreibung	Recognizes a protein of 55-62 kDa, identified as cyclin B1. In mammals, cyclin B associates with inactive p34cdc2, which facilitates phosphorylation of p34cdc2 at aa 14Thr and 15Tyr. This maintains the inactive state until the end of G2-phase. The in...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[CCNB1/2978R]
Molekulargewicht	55-62 kDa

UniProt	<a href="#">P14635</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry