

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin, pan (Epithelial Marker)(PCK/3150), CF405S conjugate, 0.1mg/mL, Clone: [PCK/3150], Mouse, Monoclonal BOT-BNC043150-100**

Artikelname	Cytokeratin, pan (Epithelial Marker)(PCK/3150), CF405S conjugate, 0.1mg/mL, Clone: [PCK/3150], Mouse, Monoclonal
Artikelnummer	BOT-BNC043150-100
Hersteller Artikelnummer	BNC043150-100
Alternativnummer	BOT-BNC043150-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC
Spezies Reaktivität	Canine, Feline, Human
Immunogen	Crude cytokeratin extract prepared from RT-4 and MCF-7 cells.
Konjugation	CF405S
Produktbeschreibung	Anti-cytokeratin clone PCK/3150 demonstrates a broad spectrum of cytokeratin reactivity. In normal tissues, PCK/3150 is reactive with most epithelial types, including bile ducts and hepatocytes in liver, bladder epithelium, breast ducts, bronchial ep...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[PCK/3150]

Molekulargewicht	40-67 kDa (Multiple)
UniProt	Multiple
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry