

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p53 Tumor Suppressor Protein (TP53/2092R), CF640R conjugate, 0.1mg/mL, Clone: [TP53/2092R], Rabbit, Monoclonal BOT-BNC402092-100

Artikelname	p53 Tumor Suppressor Protein (TP53/2092R), CF640R conjugate, 0.1mg/mL, Clone: [TP53/2092R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC402092-100
Hersteller Artikelnummer	BNC402092-100
Alternativnummer	BOT-BNC402092-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Canine, Gallus, Hamster, Human, Monkey
Immunogen	Recombinant full-length human TP53 protein
Konjugation	CF640R
Produktbeschreibung	This MAb reacts with an N-terminal epitope (aa 16-25) of both wild type and mutated p53. Mutation and/or allelic loss of p53 is one of the causes of a variety of mesenchymal and epithelial tumors. If it occurs in the germ line, such tumors run in fam...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[TP53/2092R]

Molekulargewicht	53 kDa
UniProt	P04637
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with mouse or rat Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user