

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

PTEN (Tumor Suppressor Oncoprotein) (PTEN/2110), CF640R conjugate, 0.1mg/mL, Clone: [PTEN/2110], Mouse, Monoclonal BOT-BNC402110-100

| | |
|--------------------------|---|
| Artikelname | PTEN (Tumor Suppressor Oncoprotein) (PTEN/2110), CF640R conjugate, 0.1mg/mL, Clone: [PTEN/2110], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC402110-100 |
| Hersteller Artikelnummer | BNC402110-100 |
| Alternativnummer | BOT-BNC402110-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human PTEN protein |
| Konjugation | CF640R |
| Produktbeschreibung | PTEN (phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase) contains a tensin like domain as well as a catalytic domain similar to that of the dual specificity protein tyrosine phosphatases. It was identified as a tumor suppressor that is mutated i... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [PTEN/2110] |

| | |
|------------------------|---|
| Molekulargewicht | 55 kDa |
| UniProt | P60484 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |